



International  
Animal Health Products  
THE AUSTRALASIAN COMPANY

# FARNAM<sup>®</sup> BUFFERED IODINE<sup>™</sup> SPRAY

*AN AID IN THE TREATMENT OF TOPICAL, BACTERIAL AND FUNGAL INFECTIONS IN ANIMALS.*

**PRESENTATION:** Brown liquid with alcoholic odour.

**ACTIVE CONSTITUENTS:** Each L contains:

POVIDONE IODINE 50 g/L equivalent to 5 g/L available iodine

## PROPERTIES:

About 70 % of cuts and wounds occur on the lower limb of animals, which usually heal considerably slower than skin wounds on the upper body, often delayed by contamination and infection. Lacerations, wounds, skin abrasions and low grade bacterial and fungal infectious of the skin are the most common problems of domestic and other animals.

**FARNAM<sup>®</sup> BUFFERED IODINE<sup>™</sup> SPRAY** is a safe and effective topical spray suitable for general antiseptic purposes on the skin of the most animals. It can be used to control topical infection of the skin, keeping wounds dry and covered with a protective residue film of iodine to provide prolonged activity. **BUFFERED IODINE<sup>™</sup> SPRAY** may be used in numerous surgical procedures and as a topical agent for various dermal and mucosal infections such as wounds, cuts, abrasions, burns, ringworm, navel disinfection in newly born animals, castration and docking wounds, udder wounds and cut teats.

**BUFFERED IODINE<sup>™</sup> SPRAY** contains “safe” povidone iodine that does not irritate skin or wound surface if used as recommended. **BUFFERED IODINE<sup>™</sup> SPRAY** in the quiet operating pump action pack is ideal for spray on the horses and other animals that often become frightened by the ‘hissing’ sound of pressure-pack topical sprays.

## DIRECTIONS FOR USE:

**DO NOT APPLY TO EYES OR MUCOUS MEMBRANES.**

**If used on milking cows post milking, ensure teats and udders are CLEAN and DRY before next milking to reduce residues in the milk. NOT FOR USE as a pre-milking treatment.**

**Repeated use on open wounds is not recommended.**

Shake bottle and hold upright. Apply **BUFFERED IODINE<sup>™</sup> SPRAY** using pump action. Hold bottle about 15 cm from area to be treated. Spray the affected area twice daily until healed.

**MEAT WITHHOLDING PERIOD: NIL.**

**TRADE ADVICE:** EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI contact the manufacturer on (02) 9672 7944 before using this product.

**POISONS SCHEDULE:** NOT APPLICABLE.

**REGULATORY STATUS:** AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY.

**PACK SIZES:** 500 mL spray pack.

|   |
|---|
| Australia-Wide Toll Free: 1800 801 201                        |
| Email: <a href="mailto:info@iahp.com.au">info@iahp.com.au</a> |
| Website: <a href="http://www.iahp.com.au">www.iahp.com.au</a> |