



**International**  
Animal Health Products

## CTC-ECO™ 200 GRANULAR

***FEED SUPPLEMENT FOR PROPHYLACTIC USE WHEN AT RISK FROM DISEASES AND FOR TREATMENT OF DISEASES CAUSED BY MICROORGANISMS SUSCEPTIBLE TO CHLORTETRACYCLINE IN CALVES, PIGS AND POULTRY.***

**PRESENTATION:** Brown free flowing micro-granules.

**ACTIVE CONSTITUENTS:** Each kg contains:

200 g/kg CHLORTETRACYCLINE HYDROCHLORIDE (as calcium complex)

**PROPERTIES:** CTC-ECO™ **BASE 200** is the feed stable, calcium complex of chlortetracycline hydrochloride, a broad-spectrum antibiotic, highly effective against a variety of gram-positive and gram-negative pathogenic bacteria such as Mycoplasmas, Ehrlichia, Riskettsias, Chlamidias and some atypical mycobacteria and amoeba that are not controlled by other antibiotics and bacteriostatic agents.

**CTC-ECO™ 200 GRANULAR** is specifically designed for the inclusion in animal feed. It is non-dusty micro-granules to give maximum dispersion in feed and in addition is well absorbed. It is generally very stable in the dry form, particularly when compared to other antibiotics.

### **DOSAGE AND ADMINISTRATION:**

**DO NOT USE in lactating cows where milk or milk products may be used for human consumption.** ***Prophylactic dose:*** If calves, pigs or poultry at risk of stress such as rehousing, transport, vaccination, weaning or in conditions that predispose towards disease, **CTC-ECO™ 200 GRANULAR** should be supplied at prophylactic dose for periods up to 28 days. Thoroughly mix 500 g **CTC-ECO™ 200 GRANULAR** into 1 tonne of finished feed (equivalent to 100 ppm chlortetracycline hydrochloride).

***Therapeutic dose:*** Thoroughly mix 2 kg **CTC-ECO™ 200 GRANULAR** into 1 tonne of finished feed (equivalent to 400 ppm chlortetracycline hydrochloride) and dose for not more than 7 days.

**Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease, or target species may require extending the approved withholding period.**

### **WITHHOLDING PERIOD:**

***Prophylactic dose:*** **MEAT: REMOVE ALL MEDICATED FEED, poultry – 2 days; calves & pigs – 5 days before slaughter for human consumption.**

***Therapeutic dose:*** **MEAT: REMOVE ALL MEDICATED FEED, calves – 10 days; poultry & pigs – 7 days before slaughter for human consumption.**

**EGGS – NIL**

**DO NOT USE in lactating cows where milk or milk products may be used for human consumption.** **TRADE ADVICE:** EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on (02) 9672 7944 before using this product.

**POISONS SCHEDULE:** S4 (Prescription Animal Remedy)

**REGULATORY STATUS:** AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY

**PACK SIZE:** 25 kg

Australia-Wide Toll Free: 1800 801 201

Email: [info@iahp.com.au](mailto:info@iahp.com.au)

Website: [www.iahp.com.au](http://www.iahp.com.au)

® The Australian/World symbol is a registered trademark of International Animal Health Products P/L.

™ CTC-ECO is a trademark of International Animal Health Products P/L.

© Copyright 2002–2024. All Rights Reserved.

04/2024