



International
Animal Health Products
THE AUSTRALASIAN COMPANY

MONECO® 400

BROAD SPECTRUM FEED ADDITIVE POWDER FOR CATTLE, GOATS AND POULTRY

PRESENTATION: A brown-grey free-flowing powder.

ACTIVE CONSTITUENTS: Each kg contains: 400 g MONENSIN (as MONENSIN SODIUM).

PROPERTIES: MONECO® 400 contains monensin sodium, which aids in prevention of coccidiosis in cattle (caused by *Eimeria zuerni* and *E. bovis*), goats, broiler and layer replacement chickens (caused by *E. acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix* and *E. tenella*). MONECO® 400 may be used in replacement pullets for periods up to 7 days before producing eggs and eggs products for human consumption, or as directed by a Veterinarian. With the development of intensive methods of husbandry, coccidiosis control has become more difficult. The high contamination rates and risk of infection have led to the wide use of prophylactic medication that can be maintained through MONECO® 400, especially in poultry production. In ruminants, monensin sodium has a myriad of effects on improved feed efficiency in beef cattle, weight gain, feed efficiency and reproductive performance in heifers, reduction the severity of non-clinical ketosis and by increasing energy and nutrient utilisation on higher milk production in lactating dairy cows, and is beneficial in the control of bloat in ruminants. In addition, MONECO® 400 can be used in complete feed or supplementary feed for cattle, or in supplementary feed for grazing cattle.

DOSAGE AND ADMINISTRATION: FOR ORAL ADMINISTRATION IN FEED. THE REQUIRED AMOUNT OF MONECO® 400 MUST BE THOROUGHLY MIXED IN FEED BEFORE USE ACCORDING TO THE INSTRUCTIONS.

CLAIM	COMPLETE FEEDS	SUPPLEMENTS
CATTLE: Improved feed efficiency in feedlot beef cattle except in lactating dairy cows.	Monensin 11mg - 33 mg/kg (ppm**) in ration (equiv. to 28g - 83g MONECO 400/tonne of final feed).	Monensin 50mg - 360 mg/head/day (equiv. to 125mg - 900mg MONECO 400 /head/day).
CATTLE: Aid in prevention of coccidiosis in cattle, except in lactating dairy cows.	Monensin 11mg - 33 mg/kg (ppm**) in ration (equiv. to 28g - 83g MONECO 400/tonne of final feed).	
CATTLE: Aid in control of bloat, except in dairy cows.	Monensin 25mg - 33 mg/kg (ppm**) in ration (equiv. to 63g - 83g MONECO 400/tonne of final feed).	
CATTLE: Weight gain, feed efficiency, reproductive performance (heifers).	Monensin 11mg - 33 mg/kg (ppm**) in ration (equiv. to 28g - 83g MONECO 400/tonne of final feed).	Monensin 200mg - 360 mg/head/day (equiv. to 500-900mg MONECO 400/head/day).
DAIRY COWS: Increased milk production (under some conditions) and aids reduction of severity of non-clinical ketosis incidence in lactating dairy cows.	Monensin 6mg - 18 mg/kg (ppm**) in ration (equiv. to 15g - 45g MONECO 400/tonne of final feed).	Monensin 200mg - 450 mg/head/day (equiv. to 500mg - 1,125 mg MONECO 400 /head/day).
DAIRY COWS: Aid in prevention of coccidiosis.	Monensin 11mg - 18 mg/kg (ppm**) in ration (equiv. to 28g - 45g MONECO 400/tonne of final feed).	
DAIRY COWS: Aid in control of bloat.	Monensin 11mg - 18 mg/kg (ppm**) in ration (equiv. to 28g - 45g MONECO 400/tonne of final feed).	Monensin 200mg - 450 mg/head/day (equiv. to 500mg - 1,125 mg MONECO 400 /head/day).
GOATS: Aid in prevention of coccidiosis.	Monensin 5mg - 20 mg/kg (ppm**) in ration (equiv. to 13g - 50g MONECO 400/tonne of final feed).	Monensin 7.5mg - 40 mg/head/day (equiv. to 19mg - 100 mg MONECO 400 /head/day).
POULTRY (Broilers & replacement Layers): Aid in prevention of coccidiosis.	Monensin 100mg - 120 mg/kg (ppm**) in ration (equiv. to 250g - 300g MONECO 400/tonne of final feed).	
CALVES (50-250 kg): Aid in prevention of coccidiosis.	Monensin 11mg - 33 mg/kg (ppm**) in ration (equiv. to 28g - 83g MONECO 400/tonne of final feed).	Monensin 0.5mg - 1.0 mg/kg body weight/day (equiv. to 1.3mg - 2.5 mg MONECO 400 /kg body weight/day).

** 1 ppm (parts per million) = 1 mg/kg.

PRECAUTIONS: Feeding MONECO® 400 to lactating cows may result in decreased milk fat percentage but there is no effect on total fat yield. Tiamulin (Tiamutin or Dynamutlin) should not be administered in poultry when MONECO 400 is being fed in the ration.

CONTRAINDICATIONS: The recommended levels of MONECO® 400 must not be exceeded. Horses, other equines or dogs should not be fed or allowed access to feed containing MONECO® 400.

WITHHOLDING PERIODS: (CATTLE:) MEAT and MILK: NIL. (GOATS:) MEAT: NIL; MILK: DO NOT USE in goats producing milk for human consumption. (POULTRY:) MEAT: NIL; EGGS: DO NOT USE in birds producing eggs or may in future product eggs or egg products for human consumption.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on (02) 9672 7944 before using this product.

POISONS SCHEDULE: S4 (PRESCRIPTION ANIMAL REMEDY).

REGULATORY STATUS: Australia: AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY

PACK SIZE: 20 kg and 25 kg.

Australia-Wide Toll Free: 1800 801 201

Email: info@iahp.com.au

Website: www.iahp.com.au