



International
Animal Health Products

SODIDE™ INJECTION FOR CATTLE AND HORSES

For the treatment of Actinobacillosis (woody tongue) in cattle and as an aid in treatment of Actinobacillosis in horses.

It also assists in treating infections caused by Actinomycosis (lumpy jaw), Dermatomycosis (Trichophyton spp.), Sporotrichosis and Granulomatous rhinitis in cattle and horses

PRESENTATION: Solution (injectable): 100mL, 250 mL

ACTIVE CONSTITUENTS: Each mL contains: 500 mg/ml Sodium Iodide

PROPERTIES:

SODIDE™ is used for the treatment of Actinobacillosis (woody tongue) in cattle and as an aid in treatment of Actinobacillosis in horses.

It also assists in treating infections caused by Actinomycosis (lumpy jaw), Dermatomycosis (Trichophyton spp.), Sporotrichosis and Granulomatous rhinitis in cattle and horses.

DOSAGE AND ADMINISTRATION: This product is contraindicated for use in hyperthyroid animals. If toxic signs such as excessive lacrimation, dyspnoea, tachycardia or exfoliation of the skin occur treatment should be discontinued immediately.

Not for use in pregnant mares. Use with caution in horses below 1 year of age.

This product may be given intravenously only by, or under supervision of, a veterinary surgeon.

Cattle: 40 mL/300 kg bodyweight diluted in 60 mL Water for Injections BP. This product may be given intravenously only by, or under the supervision of, a veterinary surgeon. By slow intravenous infusion, or subcutaneous injection, taking care to avoid perivascular leakage. Treatment may be repeated twice if required at 10-day intervals.

Horses: 10 mL/100 kg bodyweight by slow intravenous injection using a 14-gauge catheter taking care to avoid perivascular leakage. Can be used undiluted or diluted with sterile saline or Water for Injections BP. Treatment may be repeated twice if required at 10-day intervals.

WITHHOLDING PERIOD: **Meat**(Cattle): DO NOT USE less than 28 days before slaughter for human consumption. **Meat**(HORSES): Do not use less than 60 days before slaughter.

Milk: Milk collected 10 days following treatment MUST NOT BE USED for human consumption or processing. This milk should not be fed to bobby calves.

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI contact the manufacturer on (02) 9672 7944 before using this product.

POISONS SCHEDULE: Nil.

REGULATORY STATUS: AUSTRALIAN PESTICIDES & VETERINARY MEDICINES AUTHORITY (APVMA). APVMA Approval No. 46411/146939

IN-USE BROACHING: Discard unused portion after three months of first broaching container.

PACK SIZE: 100mL, 250ml

Australia-Wide Toll Free: 1800 801 201

Email: info@iahp.com.au

Website: www.iahp.com.au

® The Australia/World symbol is a registered trademark of International Animal Health Products P/L.

™ SODIDE is a trademark of International Animal Health Products P/L.

© Copyright 1997- 2025. All Rights Reserved.